

Nonequilibrium Fluctuations in Sedimenting Suspensions: A Dynamical Renormalization Group Theory

Alex Levine¹, Sriram Ramaswamy², and Robijn Bruinsma¹

¹*Department of Physics, University of California, Los Angeles CA 90095, USA*

²*Department of Physics, Indian Institute of Science, Bangalore 560 012, India*
(February 1, 2008)

A nonlinear two-fluid stochastic hydrodynamical description of velocity and concentration fluctuations in sedimenting suspensions is constructed, and analyzed using self-consistent (SC) and renormalization group (RG) methods. The advection of particles by velocity fluctuations is shown to be *relevant* in all dimensions $d < 6$. Both RG and SC analyses predict a strong reduction in the dependence of velocity fluctuations on system-size L relative to the $L^{1/2}$ obtained in the linearized theory of Caffisch and Luke [Phys. Fluids **28**, 785 (1985)]. This is an important step towards resolving a ten-year old puzzle in the field.

Sedimentation [1] is at once a rich and complex phenomenon in colloid science [2,3] and a frontier problem in nonequilibrium statistical mechanics. The average sedimentation speed v_{sed} is determined by balancing the driving force (gravity) against the dissipative force (viscous drag). Despite the long-ranged ($1/r$) nature of the hydrodynamic interaction between the colloids, an adequate mean-field theory [4] of v_{sed} is available for low concentrations. There is, however, no satisfactory theory of the *fluctuations* of the velocity and concentration fields in a steadily settling many-particle suspension, even at zero Reynolds number. Experiments [5–8] show clear nonthermal fluctuations in the particle motion, with the character of random stirring on a scale substantially larger than the particle size. The fluctuations δv in the settling speed are of the same order as the mean v_{sed} . Caffisch and Luke (CL) [9] showed in a *linearized* hydrodynamic analysis that, for a sedimenting system of linear dimension L , the assumption of purely *random* local concentration fluctuations led to long-ranged velocity fluctuations with $\delta v \sim L^{1/2}$. Experiments, however, find *no* dependence of δv on L [6,7,10]. In this letter we *assume* a source of randomness at small scales, the origin of which we comment on briefly below, and examine its consequences at large scales. In detail, we use a dynamical renormalization group (DRG), general scaling arguments, as well as perturbative self-consistent methods to analyze the advection of concentration fluctuations in steady-state sedimentation by the velocity fluctuations which they themselves produce. The DRG method (see Ma and Mazenko [11]), was applied to the closely related problem of the randomly stirred fluid by Forster, Nelson, and Stephen (FNS) [12]. Scaling and self-consistent approaches give a family of possible results, the simplest and most physically appealing of which accounts for the experimentally observed lack of size-dependence. The DRG results,

which are to first order in $\epsilon = 6 - d$ where d is the dimension of space, present a clear trend towards suppressing the size dependence, but simply setting $\epsilon = 3$ in these first-order calculations leaves a residual size dependence, albeit much weaker than the $L^{1/2}$ of CL [9].

We coarse-grain the dynamics on scales comparable to the random-stirring scale ℓ_s . This obliges us to use a *stochastic* description with a noise source and an effective bare diffusivity or hydrodynamic dispersion coefficient [13] $D_0 \sim v_{\text{sed}} \ell_s$. We then calculate quantities such as the velocity fluctuation on scales $\gg \ell_s$ using the following two-fluid hydrodynamic description of sedimentation along $\hat{z}$. Our coupled equations for fluctuations $\mathbf{v}$ and c in the velocity and concentration fields respectively, at zero Reynolds number, are

$$\frac{\partial c}{\partial t} + \lambda \mathbf{v} \cdot \nabla c = D_0 \nabla^2 c + \theta(\mathbf{r}, t) \quad (1)$$

$$\eta \nabla^2 v_i = m_R g \delta c P_{iz} \quad (2)$$

Here the pressure field has been eliminated by imposing incompressibility via the transverse projector $P_{ij} = \delta_{ij} - \nabla_i \nabla_j (\nabla^2)^{-1}$. Equation 1 is the advection-diffusion equation for a scalar field with $\lambda = 1$. The bare collective diffusivity D_0 and the Gaussian noise term θ , with zero mean and

$$\langle \theta(\mathbf{r}, t) \theta(\mathbf{0}, 0) \rangle = N_o \nabla^2 \delta(\mathbf{r}) \delta(t) \quad (3)$$

have the same origin — in general a combination of thermal fluctuations and hydrodynamic dispersion. If the noise is of purely chaotic origin, as in Ladd’s athermal simulation [10], then N_o and D_0 are parameters which must be determined either experimentally or by the analysis of numerical simulations. For purely thermal noise, the ratio N_o/D_0 is fixed by the fluctuation dissipation theorem (FD) [14] while $D_0 \sim k_B T / \eta a$ with η the solvent viscosity and a the radius of the colloids. Eq. 2, which expresses the balance between the driving by gravity and dissipation by viscosity, describes how concentration fluctuations produce velocity fluctuations with $m_R g$ the buoyancy-reduced colloidal weight. In Eqs. 1–2 other possible nonlinear terms, e.g. those arising from concentration-dependent mobilities, are readily shown to be subdominant relative to the advective nonlinearity $\nabla \cdot \mathbf{v} c$ [15].

If we neglect the nonlinear term (i.e. set $\lambda = 0$), then we can solve the equations to find the result of [9]:

$$\langle v_z^2 \rangle_L = 3 \langle \mathbf{v}_\perp^2 \rangle_L \propto \frac{N_o}{D_o} L \quad (4)$$

The nonlinear term causes these induced velocity fluctuations to advect further the concentration field. Since flow advection is well known to distort and suppress fluctuations [16], the ultimate size-dependence should be weaker than that predicted by Eq. 4.




FIG. 1. The renormalization group flow below the upper critical dimension in the coupling constant (λ), noise anisotropy (Γ_N) plane. The IR stable fixed point which violates the fluctuation - dissipation theorem (Γ_N^*, λ^*) is represented by the black dot at the intersection of the dashed lines representing the fixed point values of λ and Γ_N .

Let us summarize first the simple scaling and self-consistent arguments. Note first that within the above linear treatment the dynamic exponent relating spatial and temporal rescaling is $z = 2$ and the structure factor for concentration fluctuations $S(k) = \langle |c_k|^2 \rangle \neq 0$ for wavenumber $k \rightarrow 0$. The nonlinearity $\nabla \cdot \mathbf{c} \mathbf{v}$, whose advective form guarantees that there will be no graphical corrections to it under renormalization, alters these conclusions dramatically. Rescaling (isotropically for simplicity) $r \rightarrow br$, $t \rightarrow b^z t$, $c \rightarrow b^{\chi_c} c$, $v \rightarrow b^{\chi_v} v$, normalizing to retain the form of the first term in (1), and insisting that the coefficient of the nonlinearity remain unchanged, yields $\chi_c + z + 1 = \chi_v + z - 1 = 0$. We can identify the upper critical dimension d_c above which the nonlinearity is irrelevant by noting that for $d \geq d_c$, linearized behaviour should obtain: $\chi_c = -d/2$, $z = 2$. This gives $d_c = 6$, which is some indication of why a reliable answer in three dimensions is going to be hard to find. The same results can be obtained from a one-loop self-consistent calculation of the renormalized diffusivity $D(k, \omega)$ and noise spectrum $N(k, \omega)$ [15]. In either case, if we *assume* that there is no singular renormalization (except possibly logarithms) of the noise spectrum [18], we get, for $d < 6$,

$z = d/3$, and $S(k) \sim 1/D(k) \sim k^{2-d/3}$. In three dimensions this is equivalent to the result $S(k) \sim k$ of [19]. While this last is but one of a family of possible results that arise out of the scaling analysis, it has a particular physical appeal, which we discuss at the end of this Letter.

We turn now to the DRG calculation [11,12] of the renormalization of the dynamical parameters N , D , and λ . The RG recursion relations to one loop order near the upper critical dimension $d = 6$ are

$$\frac{d\bar{\lambda}}{d\ell} = \bar{\lambda}(\ell) \times \left[\frac{1}{2}\epsilon - 10^{-4}\bar{\lambda}^2(\ell) \{ 1.0 + 9.4\Gamma_N(\ell) - 3\Gamma_N^2(\ell) \} \right] \quad (5)$$

$$\frac{d\Gamma_N}{d\ell} = 2.62 \times 10^{-6}\bar{\lambda}^2(\ell) \times [1 - 5.6\Gamma_N(\ell) - 163.8\Gamma_N^2(\ell) - 231\Gamma_N^3(\ell)] \quad (6)$$

$$\frac{dN}{d\ell} = -10^{-4}N(\ell)\bar{\lambda}^2(\ell) \times [0.55 + 5.5\Gamma_N(\ell) - 5.5\Gamma_N^2(\ell)] \quad (7)$$

$$\frac{d\Gamma_D}{d\ell} = -6.72 \times 10^{-5}\bar{\lambda}^2(\ell)\Gamma_D(\ell) \times [1 + 9.63\Gamma_N(\ell)] \quad (8)$$

We have allowed explicitly for different noise strengths N_z and $N_\perp$ for wavevectors along and normal to $\hat{\mathbf{z}}$ respectively. Here, the dimensionless coupling constant is $\bar{\lambda} = \frac{\lambda N_z^{1/2}}{D_z^{1/2}}$, and $\epsilon = d - 6$, while $\Gamma_N \equiv N_\perp/N_z$ and $\Gamma_D \equiv D_\perp/D_z$ are the anisotropy factors of respectively the noise and diffusion constants. We selected the dynamical exponent such that D_z has trivial scaling. Figs. and show the resulting RG flow lines for $d > 6$ and $d < 6$ respectively. For $d > 6$, the nonlinear coupling constant $\bar{\lambda}(\ell)$ scales to zero for large ℓ . Linearized hydrodynamics is applicable and $\langle v^2 \rangle_L \propto L^{4-d} + \text{const}$ in agreement with Eq. 4. For $d < 6$, on the other hand, we find a stable fixed point $(\bar{\lambda}^*, \Gamma_N^*)$. The noise is highly anisotropic at the fixed point ($\Gamma_N^* \approx 0.0608$), while $\bar{\lambda}^* \propto \sqrt{\epsilon}$. The noise amplitude $N(\ell)$ scales to zero according to Eq. 7, as does the diffusion constant anisotropy (Eq. 8). This means that the ratio $N(\ell)/D(\ell)$ of the noise to the diffusion constant scales to zero at the fixed point signalling the absence of an FD theorem at large lengthscales below $d = 6$, irrespective of the origin of the noise. There is a FDT respecting fixed point which is unstable to any bare deviation of the noise from its FDT determined value. The linear theory, which in effect assumes an FD theorem, thus overestimates the magnitude of concentration fluctuations at large lengthscales. This flow-induced noise-suppression is highly anisotropic. This is already clear from the fact that Γ_N^* is a small number, so concentration fluctuations with wavevectors along the sedimentation direction are much larger than concentration fluctuations perpendicular to the sedimentation di-

rection. A similar conclusion can be drawn from the dynamical scaling exponents. The relaxation rate $\omega(q_z)$ for concentration fluctuations *along* the $\hat{z}$ direction depends on the wavevector q_z as $\omega(q_z) \propto q_z^{z_{\parallel}}$, with $z_{\parallel}$ the corresponding scaling exponent, while in the perpendicular direction $\omega(\mathbf{q}_{\perp}) \propto |\mathbf{q}_{\perp}|^{z_{\perp}}$. If we use $\epsilon = 3$ in Eq. 5 we find $z_{\parallel} \cong 0.91$ while $z_{\perp} \cong 1.93$. In-plane relaxation of concentration fluctuations is thus nearly purely diffusive while out-of-plane relaxation is much faster than diffusive.




FIG. 2. The renormalization group flow above the upper critical dimension in the coupling constant (λ), noise anisotropy (Γ_N) plane. The IR stable fixed point is that describing the free theory with a finite shift in the anisotropy parameter $\Gamma_N = N_{\perp}/N_z$. This fixed point model obeys the fluctuation - dissipation theorem.

To make quantitative predictions we use the dynamical scaling relations

$$N_{z,\perp}(q_{z,\perp}, \omega) = q_{z,\perp}^{\alpha} F_{z,\perp}(\omega/q_{z,\perp}^{z_{\parallel,\perp}}) \quad (9)$$

$$D_{z,\perp}(q_{z,\perp}, \omega) = q_{z,\perp}^{\beta_{z,\perp}} G_{z,\perp}(\omega/q_{z,\perp}^{z_{\parallel,\perp}}) \quad (10)$$

where $\alpha = -2 + z_{\parallel} + \gamma$, $\beta_z = -2 + z_{\parallel}$, $\beta_{\perp} = -2 + z_{\perp}$ with $\gamma = 0.83$ for $\epsilon = 3$. The scaling functions $F_{z,\perp}(x)$ and $G_{z,\perp}(x)$ are all finite at $x = 0$ but for large x , they behave as power laws:

$$\lim_{x \rightarrow \infty} F_{z,\perp}(x) \propto x^{\alpha/z_{\parallel,\perp}} \quad (11)$$

$$\lim_{x \rightarrow \infty} G_{z,\perp}(x) \propto x^{\beta_{z,\perp}/z_{\parallel,\perp}} \quad (12)$$

The full velocity fluctuation spectrum is then :

$$\langle |v_z^2(\mathbf{q}, \omega)| \rangle = \Delta \frac{q_{\perp}^4}{q^8} \left[\frac{N_z(\mathbf{q}, \omega) q_z^2 + N_{\perp}(\mathbf{q}, \omega) q_{\perp}^2}{\omega^2 + (D_z(\mathbf{q}, \omega) q_z^2 + D_{\perp}(\mathbf{q}, \omega) q_{\perp}^2)^2} \right] \quad (13)$$

$$\langle |\mathbf{v}_{\perp}^2(\mathbf{q}, \omega)| \rangle = \Delta \frac{q_{\perp}^2 q_z^2}{q^8} \left[\frac{N_z(\mathbf{q}, \omega) q_z^2 + N_{\perp}(\mathbf{q}, \omega) q_{\perp}^2}{\omega^2 + (D_z(\mathbf{q}, \omega) q_z^2 + D_{\perp}(\mathbf{q}, \omega) q_{\perp}^2)^2} \right] \quad (14)$$

with $\Delta = (m_R g / \eta)^2$. Integrating over $\mathbf{q}$ and ω while using the dynamical scaling relations given by Eqs. 9,10 gives our final result for the size-dependence of the velocity fluctuations in $d = 3$.

$$\langle |v_z|^2 \rangle_L \propto L^{1+(z_{\perp}-z_{\parallel})/2-\gamma} \quad (15)$$

$$\langle |\mathbf{v}_{\perp}|^2 \rangle_L \propto L^{1-\gamma} \quad (16)$$

The trend towards a substantial reduction of the size-dependence is gratifying. If we simply extrapolate our first-order in ϵ results to $d = 3$ we find $\langle |v_z|^2 \rangle_L \propto L^{0.6}$ and $\langle |\mathbf{v}_{\perp}|^2 \rangle_L \propto L^{0.17}$. Comparing Eqs. 15 and 16 with experiment we first note that velocity fluctuations diverge considerably more weakly than linearly along the z direction and hardly at all along the perpendicular direction so Eqs. 15 and 16 are closer to the observed size dependence than Eqs. 4 and 4. While experiments report no size dependence at all, our theory really is only reliable near $d = 6$ and a naive extrapolation of a one loop DRG calculation to $d = 3$ is questionable. Another interesting method of comparison between experiment and the present theory is in the anisotropy factor $\langle |\mathbf{v}_{\perp}|^2 \rangle / \langle |v_z|^2 \rangle$. In the linearized theory this quantity is equal to $1/3$ (Eq. 4) while the non-linear theory predicts that this ratio vanishes as $L^{-0.43}$ with system size (Eqs. 15, 16). Ladd's simulation gives 0.12 for this ratio and recent experiments 0.27 and 0.26 [6,17]. All of these measurements thus suggest that the ratio is indeed less than $1/3$.

Before closing, let us note that the naïve self-consistent results in $d = 3$ can also be obtained by dimensional arguments reminiscent of those used in the theory of homogeneous isotropic turbulence [20], but with the assumption of a scale-independent *velocity* rather than dissipation. $z = 1$ or equivalently $D(k) \sim 1/k$ means that the diffusivity at a length scale k^{-1} is given by that length times a constant which must on dimensional grounds have the units of velocity. In the absence of external, thermal noise, the only candidate for this constant is v_{sed} itself. Making the reasonable guess that *all* physical quantities, in the $Pe \rightarrow \infty$, $Re \rightarrow 0$ limit, must then be built from v_{sed} and wavenumber leads immediately to $D(k) \sim v_{\text{sed}}/k$, $\delta v \sim v_{\text{sed}}$ and the mean-square displacement of particles $\sim v_{\text{sed}}^2 t^2$ upto logarithmic factors and volume-fraction dependent coefficients.

Finally, we summarize the achievements of this work. First, starting from the correct, coarse-grained, nonlinear stochastic hydrodynamic equations, which in principle are a complete description of the problem, we have shown clearly that the origin of the disagreement between theory [9] and experiments [6,7] lies in the neglect of an important nonlinearity, *viz.*, the advection of concentration fluctuations by the fluctuations in the flow. Our work thus emphasizes that both the renormalized sedimentation speed [4] and fluctuations therein should be calculated in parallel with, and not subject to assump-

tions about, the structure factor, and shows that stochastic models are a way of doing this. Second, our identification of $d = 6$ as the upper critical dimension for the problem shows that the effects of this nonlinearity are likely to be (a) very large and (b) rather difficult to calculate analytically in three dimensions. Third, our dynamic renormalization group calculations in an expansion in $\epsilon = 6 - d$ indicate a clear trend in the direction of reduced system-size dependence. Finally, with one additional assumption, we can use self-consistent and dimensional arguments which give results which imply at most a logarithmic size-dependence, and make further predictions for experiment. A reliable calculation directly in three dimensions remains the main challenge for theory.

We would like thank Maarten Rutgers and Paul Chaikin for communicating unpublished results and for useful discussions. We would also like to thank J. Brady, D. Durian, E. Frey, E. Herbolzheimer, R. Pandit, and J. Rudnick for useful discussions. S.R. thanks F. Pincus and C. Safinya and the Materials Research Laboratory, UCSB (NSF DMR93-01199 and 91-23045), as well as the ITP Biomembranes Workshop (NSF PHY94-07194) for partial support. A. L. acknowledges support by an AT&T Graduate Fellowship.

[1] See e.g. W.B. Russel, D.A. Saville and W.R. Schowalter, *Colloidal Dispersions*, Cambridge Univ Press 1989 or R. Blanc and E. Guyon, *La Recherche* **22**, 866 (1991).
[2] R.J. Hunter, *Foundations of Colloid Science*, Oxford University Press, Oxford (1966)
[3] A.K. Sood in Solid State Physics, edited by H. Ehrenreich & D. Turnbull (Academic Press, New York, 1991) Vol. 45, p 1.
[4] G.K. Batchelor, *J. Fluid Mech.* **52**, 245 (1972)
[5] E. M. Tory, M.T. Kamel, and C. F. Chan Man Fong, *Powder Technology* **73**, 219 (1992)
[6] H. Nicolai and E. Guazzelli, *Phys. Fluids* **7**, 3 (1995)
[7] J.Z. Xue, *et al.*, *Phys. Rev. Lett.* **69**, 1715 (1992)
[8] M. Rutgers, Ph.D. thesis, Princeton University 1995 (unpublished).
[9] R.E. Caflisch and J.H.C. Luke, *Phys. Fluids* **28**, 759 (1985)
[10] The simulations of A.J.C. Ladd, *Phys. Rev. Lett.* **76**, 1392 (1996) are, however, consistent with [9]. We do not attempt to resolve this discrepancy here.
[11] S.-K. Ma and G.F. Mazenko, *Phys. Rev. B* **11**, 4077 (1975).
[12] D. Foster, D. R. Nelson, and M. J. Stephen *Phys. Rev. A* **16**, 732 (1977); D. Foster, D. R. Nelson, and M. J. Stephen *Phys. Rev. Lett.* **36**, 867 (1976)
[13] J.F. Brady and G. Bossis, *Ann. Rev. fluid Mech.* **20**, 111 (1988)
[14] U. Dekker and F. Haake *Phys. Rev. A* **11**, 2043 (1975)

[15] A. Levine, S. Ramaswamy, and R. Bruinsma, in preparation
[16] A. Onuki and K. Kawasaki, *Ann. Phys. (N.Y.)* **121**, 459, (1974); S. Ramaswamy, *Phys. Rev. A* **29**, 1506 (1984); R. Bruinsma and C. Safinya, *Phys. Rev. A* **43**, 5377 (1991)
[17] P. M. Chaikin, *Bull. Am. Phys. Soc.* March 1996.
[18] A self-consistent analysis including anisotropy by E. Frey (unpublished) appears to bear out these results.
[19] D. L. Koch and E. S. G. Shaqfeh, *J. Fluid Mech.* **224**, 275 (1991)
[20] H. Tennekes and J.L. Lumley, *A First Course in Turbulence*, MIT Press, Cambridge, USA (1972).